

Merislon[®] 12 mg tablets

(BETAHISTINE MESILATE)



Useful for relieving
Dizziness



or feeling of dizziness associated with
Meniere's disease
Meniere's syndrome



or
Vertigo

- ★ Improvement of circulatory disturbances in the inner ear
- ★ Increase of blood flow in the scala media of guinea pigs with endolymphatic hydrops
- ★ Improvement of intracerebral blood flow



DOSAGE AND ADMINISTRATION

6 mg - 12 mg

3 TIMES
DAILY
after meals

The dosage may be adjusted depending on the patient's age and symptoms.



เลขทะเบียนตำรับยาที่ 1A 190/59
Prescribing information

Composition: Each tablet contains 12 mg of betahistine mesilate **Indications:** Vertigo and dizziness associated with the following diseases: Meniere's disease, Meniere's syndrome, Peripheral vertigo **Dosage and administration:** The usual adult dosage for oral use is 6-12 mg of betahistine mesilate, three times daily after meals. The dosage may be adjusted depending on the patient's age and symptoms. **Adverse reactions:** Out of 2,254 patients treated with Merislon[®] adverse reactions were reported in 26 cases (1.15%). The most common adverse reactions were nausea in 10 patients (0.4%), vomiting in 3 patients (0.1%) and skin rash in 3 patients (0.1%). **Precaution:** Careful administration; Patients with a history of peptic ulcer or with active digestive ulcer, patients with bronchial asthma, patients with pheochromocytoma **Storage:** Do not store above 30°C *Please see further information in full reference and package insert*

Reference. Merislon package insert, version Nov 2019

โปรดอ่านรายละเอียดเพิ่มเติมในเอกสารกำกับยา

ใบอนุญาตโฆษณาเลขที่ บส.1130/2564

TH-MR-FL-21-01



Further information is available on request
Eisai (Thailand) Marketing Co., Ltd.
6th Floor, GPF Witthayu Tower A,
93/1 Wireless Road, Pathumwan, Bangkok 10330
Tel: (662) 256-6296-8 Fax: (662) 256-6299